

CHAPTER 1

AN OVERVIEW OF THE LANDSCAPE AND APPROACH FOR BIOFORTIFICATION IN AFRICA

Bouis H^{1*}, Saltzman A¹, Low J², Ball A³, and N Covic⁴

*Corresponding author email: H.Bouis@cgiar.org

¹HarvestPlus, International Food Policy Research Institute, Washington, DC

²International Potato Center, Nairobi, Kenya

³HarvestPlus, International Food Policy Research Institute, Kampala, Uganda

⁴Poverty, Health, and Nutrition Division, International Food Policy Research Institute, Addis Ababa, Ethiopia



ABSTRACT

Mineral and vitamin deficiencies are a serious public health problem in Africa. The terrible consequences of these deficiencies are well known. Biofortification is the process of breeding nutrients into staple food crops. It is one cost-effective and sustainable agricultural investment that can help to reduce mineral and vitamin deficiencies, especially in the diets of the rural poor. This chapter discusses the key questions and impact pathway around which biofortification research has been oriented over the past 15 years, and sets the stage for subsequent chapters in this special issue on biofortification for the African Journal of Food, Agriculture, Nutrition, and Development.

Key words: Biofortification, Micronutrient deficiency, Agriculture, Nutrition, Micronutrient targets, HarvestPlus

INTRODUCTION

Mineral and vitamin deficiencies are a serious public health problem in Africa. In general, dietary quality in Africa is poor, with high dependence on cereal and root staples for the bulk of dietary energy consumption, particularly among the poor [1]. Low incomes and high prices for non-staple foods such as vegetables, fruits, pulses, and animal products are the major constraints to improved dietary quality.

Non-staple foods are often dense in vitamin and minerals, and bioavailability is particularly high for animal products, yet animal products are the most expensive source of dietary energy. The poor eat large amounts of food staples to acquire dietary energy and to keep from going hungry.¹ While undernourishment has been reduced significantly for many African countries since the 1990s, accelerated improvements in diet quality and nutrition interventions are needed to reduce indicators of chronic undernutrition, including micronutrient deficiencies [2]. Because a significant portion (35 percent or more) of poor household income is spent on staple foods for energy, they have little income left to purchase vegetables, fruits and animal-based protein – dietary quality [3]. A similar story can be told for the poor in Asia and Latin America as well.

The health consequences of poor dietary quality are well known – high morbidity and infant mortality rates, compromised cognitive development for children, stunting, and low economic productivity. What people eat depends on many factors, including cultural, geographical, environmental, and seasonal factors. One of the key underlying causes leading to poor dietary quality is that current food systems do not provide minerals and vitamins in sufficient quantities at affordable prices for the poor. In non-emergency situations, poverty is a major factor that limits intake of adequate, nutritious food, which must be available, accessible, and affordable to the poor. Therefore, agricultural investments and policies that improve the availability and affordability of more nutritious foods, such as biofortification, must be made an important part of the solution.

Biofortification, the process of breeding nutrients into staple food crops, is a cost-effective and sustainable agricultural investment that can help to reduce mineral and vitamin deficiencies, especially in the diets of the rural poor. Sufficient resources to implement biofortification globally for several key staple food crops became available in 2003, which coincided with the approval of the Biofortification Challenge Program by the Technical Advisory Committee of the Consultative Group on International Agricultural Research (CGIAR).

The aim of this chapter is to describe the overall landscape and approach for biofortification in Africa and preview the chapters presented in this issue. The following chapters present both the complexity involved in implementing biofortification, and the scientific evidence that biofortification works. The plant breeding and human nutrition research to date has been largely successful and continues to generate new evidence. A

¹ Hunger, not having enough to eat to meet energy requirements, differs from malnutrition, a condition that results from a person's diet having inadequate nutrients for growth and maintenance, or poor absorption of food consumed.

critical task is to scale up the production and use of biofortified crops that are now available to farmers, particularly by impoverished and malnourished rural households, and to embed biofortification as a mainstream approach to improved micronutrient adequacy in a number of institutions.

THE AFRICAN LANDSCAPE FOR ADDRESSING MINERAL AND VITAMIN DEFICIENCIES

While there is great national and regional variation in diets in sub-Saharan Africa, most are characterized by high staple food consumption, mainly cereal or root crops. Access to micronutrient-dense food sources, including animal-source protein, fruits, and vegetables, is a major challenge for many rural households. These foods are often inaccessible because of their relatively high cost, limited local availability, and distribution challenges [4].

Vitamin A, iron, and zinc deficiencies are recognized as the most severe mineral and vitamin public health problems throughout sub-Saharan Africa. Severe vitamin A deficiency among preschool children afflicts most countries, despite widespread vitamin A supplementation programs. The prevalence of vitamin A deficiency among preschool children ranges from 40% in West and Central Africa to about 25% in southern Africa [5]. Anemia affects about 40% of pregnant women and 62% of children in Africa, about half of which is estimated to be attributed to iron deficiency [6]. Anemia levels have not significantly improved over the last 20 years [7]. Data on zinc deficiency are limited, but recent estimates suggest that 24% of Africans have inadequate zinc intakes, with pregnant women and young children at the highest risk of deficiency [8].

Furthermore, half of children with vitamin and mineral deficiencies are suffering from multiple deficiencies [9]. Table 1.1 describes rates of micronutrient deficiencies in several African countries. Micronutrient deficiencies cause poor health, low cognitive development and thus low educational attainment, and decreased work capacity and earning potential, with far-reaching consequences for national socio-economic development for current and future generations.

Several options exist to combat micronutrient deficiencies, including supplementation and food-based approaches like fortification, dietary diversification, and biofortification. For children under two, breastfeeding, micronutrient powders, and nutrient-dense complementary foods can reduce the prevalence of micronutrient deficiencies.

Vitamin A supplementation is a targeted intervention that is considered to be one of the most cost-effective interventions for improving child survival [10]. Because it is associated with a reduced risk of all-cause mortality and a reduced incidence of diarrhea [11], programs to supplement vitamin A are often integrated into national health policies. Pharmaceutical doses of synthetic vitamin A, usually in the form of gelatin capsules, are provided every six months to children under five years of age. Between 1999 and 2005, the proportion of children 6-59 months of age receiving at least one high-dose vitamin A supplement increased more than fourfold. In 2012, estimated coverage rates were near 70 percent globally [12]. However, since the funding for these interventions depends



largely on donors and international NGOs, coverage varies widely from year to year in many countries.

Commercial food fortification, where trace amounts of micronutrients are added to staple foods or condiments during processing, allows people to consume recommended levels of micronutrients. This type of fortification has been particularly successful for salt iodization: 71 percent of the world's population has access to iodized salt and the number of iodine-deficient countries has decreased from 54 to 32 since 2003 [13]. Common examples of fortification include adding B vitamins, iron, folic acid and/or zinc to wheat flour, and adding vitamin A to cooking oil and sugar. Fortification is particularly effective for urban consumers, who purchase foods that have been commercially processed and fortified. Fortification is less suitable for reaching rural consumers who often do not have access to or the incomes to afford commercially produced foods. To reach those most in need, such as poor, rural households, it may be necessary to subsidize fortified foods so that the poor do not buy cheaper non-fortified alternatives.

An alternative to commercial fortification is home, point-of-use fortification systems, in which micronutrient powders or lipid-based nutrient supplements are added to food prepared in the home. Evidence of the acceptability and efficacy of home fortification is growing [14, 15, 16], but concerns remain that it is difficult and costly to implement on a large scale. Home fortification is also subject to distribution and funding limitations.

Dietary diversity is strongly and positively associated with children's nutritional status and growth, even when controlling for socioeconomic factors [17]. In the long term, dietary diversification is likely to ensure a balanced diet that includes the necessary micronutrients. In the short term, however, investments will be required in the non-staple food and livestock sectors to increase production and reverse the trend of lower staple and higher non-staple food prices, which have made it difficult for the poor to meet their mineral and vitamin requirements through diverse diets. Investments in reducing food waste in low-income countries – by improving harvest techniques, farmer education, storage facilities, and cooling chains – are also likely to increase the availability and affordability of diverse diets [18].

Special considerations are needed for infants and young children. The transition period from breast milk or formula to solid foods is often accompanied by micronutrient deficiency in many developing countries. Food-based approaches can include additions or changes to complementary feeding practices during this period, including a focus on nutrient-dense foods and the use of specially formulated micronutrient powders. Recent Multiple Indicators Surveys and Demographic Health Surveys that have included diet diversity of children 6-23 months of age and women of reproductive age indicate that the diversity of diets for most children and mothers remain low [19, 20].

Agriculture plays a role in mineral and vitamin deficiencies

Traditionally, public research and development strategies have focused on increasing agricultural productivity in staple crops to reduce chronic calorie deficiency (undernourishment). The Green Revolution prioritized the development of high-yielding varieties of major staple crops and intensifying production, increasing the total output of



staple food crops, thereby increasing their availability and affordability by reducing their prices. From the 1970s to the mid-1990s, the price of staple foods (like rice, wheat, and maize) decreased relative to the price of micronutrient-rich, non-staple foods (like vegetables, fruits, and pulses). The same level of investment for staple crops was not made in increasing the productivity of these non-staple foods. As a result, micronutrient rich foods became relatively less affordable, particularly to the poor [21, 22].

Regional and multi-lateral organizations recognize the agriculture-nutrition link

Nutrition starts with what people eat, food products from the agricultural sector. Following the food price crises of 2008 and The Lancet's first series on Maternal and Child Nutrition (2008), the link between agriculture and nutrition has garnered more attention on the international stage. For example, the Scaling Up Nutrition (SUN) movement was launched in September 2010, bringing together governments, civil society, the United Nations, donors, private businesses, and scientists in a collective action to improve nutrition. The SUN Framework for Action calls for intensifying research on biofortification as well as on improving yields of nutrient-rich foods to better address micronutrient deficiencies [23]. Fifty-five countries have now established goals to address immediate and underlying causes of malnutrition, including through agriculture. In 2012, the CGIAR initiated a research program on Agriculture for Nutrition and Health (A4NH). The Second International Conference on Nutrition (ICN2), organized by the FAO and WHO in 2014, resulted in a political commitment to address major nutrition challenges and transform food systems through coordinated public policies. At the regional level, the African Union's 2014 Malabo Declaration recommitted the member states to the Comprehensive Africa Agriculture Development Programme (CAADP) process and committed to ending hunger and improving nutrition by 2025. Biofortification is considered an important piece of the puzzle in each of these efforts to address agriculture and nutrition linkages.

THE JUSTIFICATION FOR BIOFORTIFICATION

Biofortification provides a comparatively cost-effective, sustainable, and long-term means of delivering vitamins and micronutrients to households that might otherwise not have access to, or that cannot afford to have, a fully balanced diet. Biofortified staple food crops cannot deliver as high a level of minerals and vitamins per day as supplements or industrially fortified foods, but, based on current dietary patterns, they can help by increasing the daily adequacy of micronutrient intakes among individuals throughout the lifecycle [24]. Biofortification is not expected to treat micronutrient deficiencies or eliminate them in all population groups, but contribute to increased micronutrient intake. No single intervention will solve the problem of micronutrient deficiency, but biofortification complements existing interventions (discussed above) to provide micronutrients to the most vulnerable people in a comparatively inexpensive, cost-effective, and sustainable manner [25, 26, 27, 28, 29].

Biofortification provides a feasible means of reaching malnourished populations who may have limited access to diverse diets, supplements, and commercially fortified foods. The biofortification strategy seeks to put the micronutrient-dense trait (such as for zinc, iron or vitamin A) in basic staple food crops that are being grown and consumed by



people in developing countries and that have preferred agronomic traits, such as high yield. In contrast to complementary interventions, such as fortification and supplementation that begin in urban centers, biofortified crops reach consumers in rural areas first, since most rural farming households consume what they grow. As farmers produce and market surplus biofortified staple food crops, the intervention reaches urban areas.

Unlike the continual financial outlays required for supplementation and commercial fortification programs, a one-time investment in plant breeding can yield micronutrient-rich planting materials for farmers to grow for years to come. Biofortified varieties bred for one country can be evaluated for performance in, and adapted to, other countries and their agro-ecological growing conditions, thereby potentially multiplying the benefits of the initial investment. As seed producers incorporate biofortified crops into their product lines, biofortification becomes more sustainable over time, provided that regulatory mechanisms are in place to maintain standards and related claims. While recurrent expenditures are required for monitoring and maintaining these traits in crops, these are low compared to the cost of the initial development of the nutritionally improved crops and the establishment, institutionally speaking, of nutrient content as a legitimate breeding objective for the crop development pipelines of national and international research centers. Once established, the cost of maintaining biofortified traits represents a small portion of ongoing global investment in crop improvement.

There are three common approaches to biofortification: agronomic, conventional, and transgenic. Agronomic biofortification provides temporary micronutrient increases through fertilizers and/or foliar sprays. This approach is useful if the goal is to increase micronutrients that can be directly absorbed by the plant, such as zinc, but is less efficient and effective for micronutrients that are synthesized in the plant and cannot be absorbed directly [30]. Conventional plant breeding involves identifying and developing parent lines with high vitamin or mineral levels and crossing them over several generations to produce plants with the desired nutrient and agronomic traits. Transgenic plant breeding seeks to do the same in crops where the target nutrient does not naturally exist at the required levels. The chapters in this special issue focus only on conventionally-bred biofortified varieties of staple food crops and not on transgenic plant breeding, as the evidence base has been developed with conventionally-bred biofortified crops. Transgenic biofortified crops have not been released in any African country.

UNPACKING THE STEPS TO SUCCESSFUL IMPLEMENTATION OF BIOFORTIFICATION

As with fortification, an initial question for biofortification was: what level of extra minerals and/or vitamins added to diets would be required to have a measurable public health impact? The answer is complex, depending on age- and gender-specific nutrient requirements, per capita consumption of a particular food, bioavailability of the nutrients, and nutrient retention – as shown in the equation below:



Extra Nutrient Supplied
Through Biofortification

----- = Additional Percentage of Estimated Average Requirement Supplied, where
Nutrient Requirement

Extra Nutrient Supplied By Biofortification =

Per Capita Consumption of the Food Staple	X	Increment in Density of Mineral/Vitamin Due to Plant Breeding	X	Retention of Mineral/ Vitamin in Processing/ Storage/Cooking	X	Percent Bioavailability of Mineral/Vitamin as Consumed
(Term A)		(Term B)		(Term C)		(Term D)

A simple example provides clarity:

- An adult woman in a maize-eating society may consume, on average, the equivalent of 300 grams of shelled, dried maize per day (Term A).
- White maize has zero provitamin A. The plant breeding target for biofortified, orange maize is 15 mg of provitamin A carotenoids per kilogram of shelled, dried corn at harvest, units sometimes referred to as 15 parts per million (ppm). The *increment* due to plant breeding, therefore, is +15 ppm (Term B).
- 70% of provitamin A carotenoids may be lost during processing/storage/cooking; retention is, therefore, 30% (Term C).
- After ingestion, four units of provitamin A carotenoids are converted to one unit of absorbed retinol²; bioavailability, therefore, is 25% (Term D).

Consuming biofortified maize on any given day, a one-for-one substitution for white maize, then adds:

300 grams X +15 mg/kg provitamin A carotenoids X 30% X 25% = +0.34 mg of absorbed retinol per day.

The estimated average requirement (EAR) for absorbed retinol (vitamin A) for adult women is 0.85 mg retinol per day. Therefore, biofortification would provide an extra 40% of the EAR per day.

The numbers used in the above example are realistic, based on research described in Chapters 2, 3, and 4. However, educated guesses had to be made for each of these magnitudes when plant breeding was initiated. Plant breeders, for example, concluded in 2005 that there was a reasonable probability that they could develop maize varieties with 15 ppm provitamin A carotenoids in high-yielding backgrounds, but they could not be sure. Today, some experimental maize lines, not yet in high-yielding backgrounds, have more than 30 ppm provitamin A carotenoids. Moreover, nutritionists surmised in 2005 that an additional 40 percent of the EAR for vitamin A in diets would have a positive public health impact, but, similarly, nutritionists could not be sure. Economists and nutritionists used these numbers to calculate, on an *ex-ante* basis, the cost-effectiveness

² When consumed, provitamin A carotenoids are metabolized by the body to retinol, the absorbable form of vitamin A. Bioavailability of provitamin A carotenoids is measured by retinol activity equivalents (RAE). In the case of orange maize, research demonstrated that 1 RAE was converted for each 4 mg/kg of provitamin A carotenoids consumed.



of biofortification under various farmer adoption rates in selected developing countries. The methodologies used and calculations, which showed biofortification to be highly cost-effective, are discussed in Chapters 13 and 14. These calculations provided the quantitative evidence for donors to make and sustain investments in biofortification for the ten years that it took to develop, test, and release biofortified staple food crop varieties in a significant number of developing countries.

Yet provitamin A in maize is only one example of biofortified crop success. Table 1.2: Revised Assumptions and Target Levels, shows the magnitudes used for the terms in the equation above for a number of biofortified crops in setting plant breeding targets, and for making decisions as to what crop-nutrient combinations were selected for investment and eventual deployment. Originally set using limited data on consumption patterns, as well as nutrient stability and retention in the biofortified crops, nutrient targets have been validated and updated for specific target populations as more data became available. These revised assumptions are reported in Table 1.2 and further described in Chapters 2, 3, and 4. The provitamin A levels in orange sweet potato are sufficiently high that 100 percent of the EAR is met for both target groups listed in Table 1.2. The micronutrient target levels initially set in 2005 were later increased for zinc-biofortified crops, but ongoing research showed that targets were adequate for vitamin A and iron-biofortified crops. Nutritionists continue to monitor and evaluate results and they revise assumptions and targets as new evidence emerges.

After initially setting nutrient targets, researchers identified three broad questions to be addressed in order for biofortification to be successful:

- Can breeding increase the micronutrient density in food staple crops to target levels that will have a measurable and significant positive impact on nutritional status?
- When consumed under controlled conditions, will the extra nutrients bred into the food staple crops be absorbed and utilized at sufficient levels to improve micronutrient status?
- Will farmers be willing to grow the biofortified varieties and will consumers be willing to buy and eat them in sufficient quantities?

To answer these questions, researchers carried out a series of activities classified in three phases of discovery, development, and delivery, as shown in Figure 1.1: Biofortification Impact Pathway, below. Since 2012, biofortification has moved significantly into the delivery phase in several African countries.






Figure 1.1: Biofortification Impact Pathway

Within the final steps of the impact pathway, new areas for inquiry have emerged, including the role of processors and private sector actors in the marketing of biofortified varieties. With strong proof-of-concept for biofortification, moving towards scale will require increased public and private sector investment in crop development and seed systems to sustain the pipeline of biofortified varieties. The following chapters review the evidence developed to date along the impact pathway and chart the path for future research and scaling.

ORGANIZATION OF THE CHAPTERS IN THIS VOLUME

Developing and delivering biofortified crops has required donor buy-in and investment, evidence of the potential to address the targeted micronutrient deficiencies, and promoting adoption and sustainability at the country level. For biofortification to successfully address micronutrient deficiencies, the given micronutrients must be present in sufficient amounts, adequately retained during processing and storage, and bioavailable for absorption. Chapters 2, 3, and 4 address these aspects, presenting summaries of the nutrition and food science research undertaken to date on consumption levels of staple food crops, nutrient retention, bioavailability, and efficacy. In Figure 1.1: Biofortification Impact Pathway, this research corresponds to the first two steps under discovery, and the second step under development.

Crop development requires effective screening and testing of varieties. Chapters 5 and 6 discuss the progress in crop development using conventional plant breeding methods.

Investments in crop development have first focused on the most widely consumed food staple crops, and more than 150 varieties of biofortified crops have been developed and released in more than 30 countries (Chapter 5). In Chapter 6, measurement of minerals and vitamins in the edible portions of biofortified crops and foods made from such crops are discussed, including innovations in high throughput, low-cost analytical methods. These correspond to step 3 under discovery, step 1 under development and step 1 of delivery in the Biofortification Impact Pathway.

Chapters 7-12 primarily address the third step under development in the Biofortification Impact Pathway, and both steps under delivery, but include some findings related to discovery as well. Chapters 7, 8, 9, and 10 are crop specific – discussing the range of discovery-development-delivery experiences to date in Africa for four biofortified crops – orange sweet potato across the continent, vitamin A orange maize in Zambia, vitamin A yellow cassava in Nigeria, and iron beans in Rwanda, respectively. General marketing and branding issues are discussed in Chapter 11. Integrating biofortified crops into existing international development projects is explored in Chapter 12, which focuses on the use of biofortified crops in World Vision International’s programs.

Chapters 13, 14, and 15 focus on economic analysis. Chapter 13 summarizes the Theory of Change and how the nutritional impact of biofortified crops is measured and maximized. Chapter 14 examines complementarities and tradeoffs among a range of micronutrient interventions – supplementation, fortification, and biofortification -- in three countries. This analysis is also *ex-ante*. Chapter 15, however, is a summary of *ex-post* findings of the pilot delivery of orange sweet potato to white sweet potato-growing farm households in Mozambique and Uganda – using a randomized, control testing design.

The importance of advocacy to build stakeholder and policy support for scaling up farmer adoption and the sustainable mainstreaming of biofortification is discussed in Chapter 16, including recommendations for a forward-looking advocacy strategy at the national level. Several countries where biofortified crops are available (including Rwanda, Zambia, Mozambique, and DRC) have incorporated biofortification into their national nutrition strategies. Biofortification programs are increasingly supported by national governments, particularly in China, India, and Brazil, as well as several other countries in Latin America. The specific policy approaches to incorporating biofortified crops into regional and national nutrition strategies are discussed in this chapter. Finally, Chapter 17 evaluates lessons learned from all previous chapters and charts a proposed way forward for accelerating the integration of biofortified crops in African diets.

ACKNOWLEDGEMENTS

The editors of this special issue would like to acknowledge all contributing authors and all other scientists and practitioners who continue to dedicate their efforts to developing and delivering biofortified crops.



Table 1.1: Micronutrient Deficiency Status of Select African Countries

Country	Total Pop ('000)	Proportion of pre-school age children with anemia (Hb<110 g/L)	Proportion of non-pregnant women with anemia (Hb<120 g/L)	Proportion of pre-school age children with vitamin A deficiency (serum retinol <0.70 µmol/l)	Proportion of population at risk of inadequate intake of zinc	Proportion of children <5 stunted	Total Population with Iron Deficiency (Considered 1:1 ratio with Anemia) ('000)	Total Population with Vitamin A Deficiency ('000)	Total Population with Inadequate Zinc Intake ('000)	Total population at risk of Micronutrient Deficiency with Anemia and Stunting
Nigeria	175,288	71.0	47.3	29.5	12.8	36.4	50,573	9,124	22,436	70,263
Ethiopia	90,178	49.5	18.9	46.1	21.7	44.2	12,575	6,227	19,568	24,131
Congo, Dem. Rep.	73,291	67.1	49.0	61.1	57.5	43.5	20,904	8,040	42,142	34,434
Tanzania	50,705	60.8	38.2	24.2	22.9	34.8	12,369	2,241	11,611	17,755
Uganda	37,923	56.2	25.6	27.9	23.8	33.7	7,352	2,116	9,025	11,858
Ghana	26,721	76.1	55.9	75.8	21.0	22.7	8,519	2,859	5,611	12,216
Mozambique	25,590	66.5	43.7	68.8	60.5	43.1	6,665	2,818	15,482	11,215
Malawi	16,953	65.6	27.6	59.2	34.2	47.8	3,698	1,899	5,798	7,099
Zambia	14,767	57.7	28.2	54.1	38.0	45.8	3,076	1,577	5,611	5,958
Rwanda	11,950	37.6	17.1	6.4	39.8	44.3	1,530	167	4,756	2,620

Hb: Hemoglobin

ID: iron deficiency

IZI: inadequate zinc intake

VAD: vitamin A deficiency

Source: Authors' calculations based on WHO (2009), WHO (2015), Hotz (2004), and IFPRI (2014).

Table 1.2: Revised Assumptions and Target Levels, 2016*Based on research on intake, bioavailability, and retention for biofortified crops*

Assumptions (revised)	IRON		ZINC		VITAMIN A	
Age/physiological status group	Non-pregnant, non-lactating women	Children 4-6 yr of age	Non-pregnant, non-lactating women	Children 4-6 yr of age	Non-pregnant, non-lactating women	Children 4-6 yr of age
Daily physiological requirement ($\mu\text{g}/\text{day}$) ¹	1,460 μg	500 μg	2,960 μg	1,390 μg	500 μg REA ²	275 μg REA ²
	Pearl Millet (whole grain flour) India		Wheat (whole grain flour) India		Maize (whole to maize meal) Zambia	
Intake (g/day)	222	87	258	71	287	172
Micronutrient retention after processing (%)	90		95		37	
Bioavailability (%)	7.5		15		17 (6:1)*	
Baseline micronutrient content ($\mu\text{g}/\text{g}$)	47		25		0.5	
Additional content required ($\mu\text{g}/\text{g}$)	+30		+18		+15	
Total final content ($\mu\text{g}/\text{g}$)	77		43		15.5	
% of the requirement contributed by biofortification	30	35	22	13	54	59
% of the requirement contributed by entire micronutrient content of staple (baseline + biofortification) (%)	79	90	53	31	56	61
	Beans (boiled thick broth) Rwanda		Rice (polished parboiled) Bangladesh		Cassava (fresh weight to cassava meal) Nigeria	
Intake (g/day)	198	107	422	159	940	348
Micronutrient retention after processing (%)	90		90		35	
Bioavailability (%)	7		25		20 (5:1)*	
Baseline micronutrient content ($\mu\text{g}/\text{g}$)	50		16		0	
Additional content required ($\mu\text{g}/\text{g}$)	+44		+12		+15	
Total final content ($\mu\text{g}/\text{g}$)	94		28		15.0	
% of the requirement contributed by biofortification (%)	38	59	39	31	197	133
% of the requirement contributed by entire micronutrient content of staple (baseline + biofortification) (%)	80	127	90	72	197	133

Assumptions	Vitamin A	
Age/physiological status group	Non-pregnant, non-lactating women	Children 4-6 yr of age
Estimated average requirement ($\mu\text{g}/\text{day}$) ^a	500 μg REA	275 μg REA
	Orange-fleshed sweet potato (Uganda)	
Intake (g/day)	167	101
Retention, boiled (%)	80	
Bioavailability (%)	8	
Baseline micronutrient content ($\mu\text{g}/\text{g}$)	0	
Additional content required ($\mu\text{g}/\text{g}$)	+70	
Total final content ($\mu\text{g}/\text{g}$)	70	
% of the requirement contributed by biofortification (%)	160	176
% of the requirement contributed by entire micronutrient content of staple (baseline + biofortification) (%)	160	176

*beta carotene equivalent to retinol bioconversion ratio

¹Physiological requirements are derived from the IOM 2001 Dietary Reference Intakes for iron; from EFSA Journal 2014 for zinc; and from IOM 2001 for vitamin A.

²Retinol activity equivalents

REFERENCES

1. **Food and Agriculture Organization of the United Nations (FAO)** Regional overview of food insecurity: African food security prospects brighter than ever. FAO, Accra, 2015.
2. **International Food Policy Research Institute (IFPRI)** Global Nutrition Report 2016: From promise to impact: ending malnutrition by 2030. IFPRI, Washington, DC, 2016.
3. **Food and Agriculture Organization of the United Nations (FAO)** The State of Food Insecurity in the World. FAO, Rome, 2011.
4. **Fanzo J, Hunter D, Borelli T, and F Mattei** *Diversifying Food and Diets: Using Agricultural Biodiversity to Improve Nutrition and Health*. New York, Routledge, 2012.
5. **World Health Organization (WHO)** Global Prevalence of Vitamin A Deficiency in Populations at Risk 1995–2005: WHO Global Database on Vitamin A Deficiency. WHO, Geneva, 2009.
6. **World Health Organization (WHO)** The Global Prevalence of Anaemia in 2011. WHO, Geneva, 2015.
7. **Haddad L, Bendeck MA, Bhatia K, Eriksen K, Jallow I, and N Ledlie** Africa's Progress Toward Meeting Current Nutrition Targets. In: Covic NM and S Hendriks (Eds). ReSAKSS Annual Outlooks and Trends Report Citation Annual Trends and Outlook Report 2015. International Food Policy Research Institute, Washington, DC, 2016: 12-27.
8. **Bailey R, West KP Jr. and RE Black** The Epidemiology of Global Micronutrient Deficiencies. *Annals of Nutrition & Metabolism* 2015; **66**(suppl 2): 22-33.
9. **Micronutrient Initiative and UNICEF** Vitamin and Mineral Deficiency: A Global Progress Report. Micronutrient Initiative, Ottawa, 2009.
10. **Tan-Torres Edejer T, Aikins M, Black R, Wolfson L, Hutubessy R, and D Evans** Cost Effectiveness Analysis of Strategies for Child Health in Developing Countries. *British Medical Journal* 2005; **331**: 1177.
11. **Imdad A, Herzer K, Mayo-Wilson E, Yakoob MY, and ZA Bhutta** Vitamin A Supplementation for Preventing Morbidity and Mortality in Children from 6 months to 5 Years of Age. *Cochrane Database of Systematic Reviews* 2010; **12**:CD008524.



12. **United Nations Children's Fund (UNICEF)** State of the World's Children 2014. UNICEF, New York, 2014.
13. **Andersson M, Karumbunathan V, and MB Zimmermann** Global Iodine Status in 2011 and Trends over the Past Decade. *Journal of Nutrition* 2012; **142**: 744–750.
14. **Adu-Afarwuah S, Lartey A, Brown K, Zlotkin S, Briend A and K Dewey** Home Fortification of Complementary Foods with Micronutrient Supplements is Well Accepted and Has Positive Effects on Infant Iron Status in Ghana. *American Journal of Clinical Nutrition* 2008; **87(4)**: 929–938.
15. **Dewey K, Yang Z and E Boy** Systematic Review and Meta-Analysis of Home Fortification of Complementary Foods. *Maternal and Child Nutrition* 2009; **5(4)**: 283–321.
16. **de Regil, LM, Suchdev P, Vist G, Walleser S and JP Pena-Rosas** Home Fortification of Foods with Multiple Micronutrient Powders for Health and Nutrition in Children Under Two Years of Age. *Evidence-Based Child Health* 2013; **8**: 112–201.
17. **Arimond M and MT Ruel** Dietary Diversity is Associated with Child Nutritional Status: Evidence from 11 Demographic and Health Surveys. *Journal of Nutrition* 2004; **134(10)**: 2579–2585.
18. **Food and Agriculture Organization of the United Nations (FAO)** Global Food Losses and Food Waste – Extent, Causes and Prevention. FAO, Rome, 2011.
19. **United Nations Children's Fund (UNICEF)** Statistics and Monitoring – Country Statistics. 2015. Retrieved from: http://www.unicef.org/statistics/index_countrystats.html on September 3, 2015.
20. **Demographic and Health Survey Programs (DHS)** All Surveys by Country. 2016. Retrieved from: <http://dhsprogram.com/What-We-Do/survey-search.cfm?pgtype=main&SrvyTp=country> on October 6, 2016.
21. **Bouis HE** Improving Human Nutrition Through Agriculture. *Food and Nutrition Bulletin* 2000; **21**: 549-565.
22. **Kennedy E and HE Bouis** Linkages Between Agriculture and Nutrition: Implications for Policy and Research. International Food Policy Research Institute, Washington DC, 1993.
23. **Scaling Up Nutrition (SUN)** Scaling Up Nutrition, A Framework of Action. 2011. Retrieved from: http://scalingupnutrition.org/wp-content/uploads/pdf/SUN_Framework.pdf on November 10, 2011.

24. **Bouis HE, Hotz C, McClafferty C, Meenakshi JV and WH Pfeiffer** Biofortification: A New Tool to Reduce Micronutrient Malnutrition. *Food and Nutrition Bulletin* 2011; **32 (Supplement 1)**: 31S–40S.
25. **Bouis HE** Economics of Enhanced Micronutrient Density in Food Staples. *Field Crops Res.* 1999; **60**:165–173.
26. **Nestel P, Bouis HE, Meenakshi JV and WH Pfeiffer** Biofortification of Staple Food Crops. *J. Nutr.* 2006; **136**:1064–1067.
27. **Pfeiffer WH and B McClafferty** HarvestPlus: Breeding Crops for Better Nutrition. *Crop Sci.* 2007; **47**:S88–S105.
28. **Qaim M, Stein AJ and JV Meenakshi** Economics of Biofortification. *Agric. Econ.* 2007; **37**:119–133.
29. **Meenakshi JV, Johnson N, Manyong V, DeGroot H, Javelosa J, Yanggen D and F Naher** How Cost-Effective is Biofortification in Combating Micronutrient Malnutrition? An Ex Ante Assessment. *World Development* 2010; **38 (1)**: 64–75.
30. **Lyons G and I Cakmak** Agronomic Biofortification of Food Crops with Micronutrients. In Bruulsema TW, Heffer P, Welch RM, Cakmak I and K Moran (Eds.) *Fertilizing Crops to Improve Human Health: A Scientific Review*. Paris, France: International Plant Nutrition Institute, 2012: 97-122.
31. **Hotz C and Brown KH** International Zinc Nutrition Consultative Group (IZiNCG) Technical Document #1. *Food and Nutrition Bulletin* 2004; **25 (1)**: s4.
32. **International Food Policy Research Institute (IFPRI)** Global Nutrition Report 2014: Action and accountability to accelerate the world's progress on nutrition. IFPRI, Washington, DC, 2014.
33. **Institute of Medicine** Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. Washington, DC: National Academy Press, 2001.
34. **European Food Safety Authority (EFSA)** Scientific Opinion on Dietary Reference Values for zinc. *EFSA Journal* 2014; **12 (10)**: 3844.